

RELATÓRIOS DE CAMPANHA

INTERNATIONAL SURVEY FOR THE ASSESSMENT OF THE STRENGTH OF THE
SARDINE AND ANCHOVY RECRUITMENT IN ATLANTIC IBERIAN WATERS
“IBERAS1119”



Pablo Carrera, Pedro Amorim, Ana, Morenó, María Manuel Angélico
e Paz Díaz Conde



Edição
IPMA
Rua C – Aeroporto de Lisboa
1749-007 LISBOA
Portugal

Edição Digital
Conceição Almeida

Capa
Anabela Farinha / Conceição Almeida

Disponíveis no sítio web do IPMA
<http://ipma.pt/pt/publicacoes/index.jsp>

Todos os direitos reservados

Referência Bibliográfica

CARRERA P., AMORIM P., MORENO A., ANGÉLICO M.M., CONDE P.D., 2019. International survey for the assessment of the strength of the sardine and anchovy recruitment in Atlantic Iberian waters “IBERAS1119”. *Relatórios de Campanha*, 49p. (<http://ipma.pt>)

INTERNATIONAL SURVEY FOR THE ASSESSMENT OF THE STRENGTH OF THE SARDINE AND ANCHOVY RECRUITMENT IN ATLANTIC IBERIAN WATERS



IBERAS1119 SURVEY REPORT



**INSTITUTO ESPAÑOL DE OCEANOGRAFÍA
INSTITUTO PORTUGUÊS DO MAR E DA ATMOSFERA**

INDEX

INTRODUCTION	5
OBJECTIVES	7
MATERIAL AND METHODS	7
Working Area	7
Acoustic	9
1 NASC allocation.....	9
2 Echointegration estimates.....	12
3 Centre of Gravity	12
Fishing stations	12
Plankton and hydrological characterisation.....	13
Top predator observations	13
Fish sampling.....	13
1 Catch and length distribution per specie	13
2 Weight Length relationship	13
3 Biological sampling.....	13
RESULTS.....	14
HYDROGRAPHIC CONDITIONS	14
ACOUSTIC.....	16
School extraction and total backscattering energy.....	16
Fishing station and echotrace allocation	18
1 Chub mackerel echotrace identification	18
2 Boar fish echotrace identification.....	19
3 Anchovy echotrace identification.....	20
4 Sardine fish echotrace identification	20
5 Fishing station used for echotrace allocation	20
Acoustic assessment	23
1 Sardine assessment.....	24
2 Anchovy assessment.....	32
3 Horse mackerel assessment	38
4 Chub mackerel assessment.....	41
DISCUSSION AND CONCLUSIONS.....	43
CONSULTED IBLIOGRAPHY	46

TECHNICAL SUMMARY

Institution: INSTITUTO PORTUGUÊS DO MAR E DA ATMOSFERA/INSTITUTO ESPAÑOL DE OCEANOGRAFÍA
Survey name: IBERAS1119
Vessel name: Angeles Alvariño (46.70 m length, 10.50 width 988 GRT, 900 kW diesel-electric)
Dates: 05-27/09/2019
Area: WESTERN IBERIAN COAST (9aCS-9aCN-9aN)
Type: Acoustic-Trawl
Main objective: Biomass estimation by means of echointegration of the main pelagic fish population present in the surveyed area. Physical, chemical and biological characterisation of the pelagic ecosystem.
Sampling strategy Systematic grid with random start, tracks 4/8 nmi apart from 20 to 100 isobath
Main sampling procedures EK-80 at 18-38-70-120-200 kHz acoustic frequencies. 839 nmi prospected. Only day time
Pelagic fishing stations: 16
Marine mammals and birds observations (not yet determined)
Hydrological characterisation. 48 stations

Personnel	AMORÍN, PEDRO	MAGALHÃES DA SILVA, ELISABETE
1 st leg	ANGELICO, MARIA MANUEL	MORENO MARQUES, ANA
Vigo/Lisboa	CARRERA LÓPEZ, PABLO	MURCIA ABELLÁN, JOSÉ LUIS
Dates: 07 to 21/09	DA CONCEIÇÃO MENDOZA, ANTONIO PEDRO	SÁNCHEZ BARBA, MARÍA
	DÍAZ CONDE, PAZ	SÁNCHEZ HERMOŚÍN, PABLO
	DOS SANTOS BARRA, JORGE	SANCHO MARTÍNEZ, PAULA
	MACHADO DE ABREU, PAULA	
2 nd leg	AMORÍN, PEDRO	MURCIA ABELLÁN, JOSÉ LUIS
Lisboa/Cádiz	AUTÓN DÍAZ, URBANO	PASTOR, JOÃO
Dates: 22-27/09	CARRERA LÓPEZ, PABLO	PEREIRA GAMEIRO, CARLA
	DA CONCEIÇÃO MENDOZA, ANTONIO PEDRO	PEREIRA NUNES, CRISTINA
	DÍAZ CONDE, PAZ	SÁNCHEZ BARBA, MARÍA
	FEIJÓ, DIANA	SÁNCHEZ HERMOŚÍN, PABLO
	MAGALHÃES DA SILVA, ELISABETE	

Report authors Pablo Carrera, Pedro Amorim, Ana, Moreno, María Manuel Angélico, Paz Díaz Conde

INTRODUCTION

According to ICES, the sardine biomass of age 1 and older fish has decreased since 2006; it has been below B_{lim} since 2009; and it has stabilized to a historical low since 2012. Recruitment has been below the long-term average since 2005 and in 2017 it was estimated as the lowest in the time-series. Fishing mortality has been above F_{lim} for most of the time-series but has been decreasing from a peak in 2011. In 2017, it is the lowest in the time-series and around F_{pa} . Although sardine is not considered a short-lived species, the lack of enough adults, resulted in a very low presence of older ages (e.g. very low expectation for reaching ages older than 5 due to the high natural mortality), being the bulk of the population composed by younger fish, which in turn, make this species looks like a short-lived species.

In such conditions, any recovery of the biomass will likely be triggered by the strength of the recruitment. Thus when juveniles can be assessed at age 0, the estimates can be used to predict the relative strength of the future recruitment to the fisheries. This strategy is of special interest to manage the fisheries for short-lived species because of the short time between spawning and the

exploitation of subsequent emerging recruits.

On the other hand, in coincidence with the decrease of sardine, off north Portugal and South Galicia, anchovy population has sharply increased. Monitoring this outburst is, therefore of interest as this species would partially compensate the recent lack of sardine.

Attending the experience achieved by IPMA through JUVESAR time series, an acoustic survey targeted on sardine recruitment, and by Azti and IEO in the JUVENA survey to improve the assessment and management of the Bay of Biscay anchovy together with ECOCADIZ recruit survey conducted by IEO in the Gulf of Cadiz, IBERAS will try to get a recruitment index for both species in Atlantic waters of the Iberian peninsula, aiming at to improve the estimation of the strength of the recruitment of both Iberoatlantic sardine and the western component of the south anchovy population.

In 2018 the survey was undertaken in November. However, both the bad weather conditions, that limited the number of effective survey days, and the aggregation and distribution patterns of the fish, with rather isolate and big schools (figure 1) that made difficult either to find and, specially, to improve the precision of the biomass estimates (figure 2), led to change the period of the survey. Therefore, the survey was shifted to September, at the same time of JUVENA, which in turn allows a synoptic coverage of the Iberian peninsula at the end of summer, beginning fall.




Figure 1: mega-school of anchovy recorded in 2018 during IBERAS north Figueira da Foz




Figure 1: Cumulated backscattering energy per track in IBERAS1118. 3 of them are highlighted due to contribution to the total energy.

OBJECTIVES

- i. Acoustic estimates by echointegration of the strength of the anchovy and sardine recruitment off Portugal and south Galicia
- ii. Oceanographic (physical -CTD- and biological _bongo nets) characterization of the surveyed area
- iii. Charting the relative abundance of apical predator along the surveyed area

MATERIAL AND METHODS

Survey was carried out on board R/V Angeles Alvariño, a similar vessel of Ramón Margalef, used in the previous survey IBERAS1118, from 5th until 27th September, departing from the port of Vigo and arriving to Cádiz harbour on the evening of 27th.

A scale was scheduled in Lisbon on 21st. Two first days were used to calibrate the transducers. For this purpose, the vessel moored in the Pontevedra Bay. The wind strength did not to allow to calibrate de vessel during the first day, which was completed on 6th.

Working Area

From Fisterra cape until São Vicente cape, from shoreline (20 m) to 100 m isobath over an adaptive grid with tracks distance between 4-8 nmi on account the potential recruitment distribution area of both sardine and anchovy. Tracks were enlarged or shorted accordingly. Figure 3 is showing the foreseen survey track.

The methodology was similar to that of the previous surveys and is summarised in ICES Cooperative Research Report No. 332. 268 pp. <https://doi.org/10.17895/ices.pub.4599>. The backscattering acoustic energy from marine organisms was measured continuously during daylight except in the northern area where some tracks were steamed at night. Pelagic trawls were carried out whenever possible to help identify the species (and size classes) that reflect the acoustic energy. During daylight hours, concurrently to acoustics, a trained observer recorded marine mammal, seabird, floating litter and vessel presence and abundance.

At night, when acoustics surveying is not running, CTD profiles for hydrography and zooplankton samples (Bongo 60 and Manta trawl nets) are collected, opportunistically, in some of the transects.

Besides, in specific areas chosen on the core expected distribution area of juveniles, the very shallower waters (15-10 m) were prospected with a portable EL60 with a 120 kHz transducer. For this purpose the auxiliary dinghy of the vessel was used. As shown in figure 4, the normal tracks (dotted lines) were extended towards the coast (black line), which were prospected by the dinghy. Simultaneously, the vessel steamed the intertrack line (red lines). Results at 120 kHz recorded by both echosounder (EK80 on board Angeles Alvariño and EK60 on board dinghy) were compared.




Figure 3: Survey track

Table1. Expected survey coverage and time in each ICES Sub-Division

Zone	No tracks	No of nautical miles		Acoustic	Fishing st.	TOTAL
		track	Unión	hr-days	hr-days	hr-days
Calibration						2
Plataforma 9a N	9	83	75	15-1	12-0.86	27-2
Rías Baixas (9a-N)	23	112	0	8-1	12-0.86	20-2
O. Norte (9a-CN): Caminha-Porto	6	78	48	11.8-1	12-0.86	24-2
O. Norte (9a-CN): Porto-Figueira	12	189	46	23.4-1.67	16-1.14	39.4-3
O Norte (9a-CN): Figueira-Nazaré	10	109	34	14.1-1	8-0.57	22.1-2
O Sul (9a-CS): Nazaré-Roca	9	100	59	15.9-1.14	8-0.57	24-2
O Sul (9a-CS): Roca-Troia	15	141	59	14.09-1	16-1.14	30-3
O Sul (9a-CS): Troia-São Vicente	12	81	78	15.8-1.13	16-1.14	31.8-3
Total	96	831	431	127-10.5	96-6-86	222.23-(17-19)




Figure 4: Acoustic scheme in shallower waters

Acoustic

Acoustic equipment consisted on a Simrad EK-80 scientific echosounder, operating at 18, 38, 70, 120 and 200 kHz, working in CW mode. All frequencies were calibrated according to the standard procedures (ICES-CRR326) during the first two days. The elementary distance sampling unit (EDSU) was fixed at 1 nm. Acoustic data were obtained only during daytime at a survey speed of 8-10 knots, although, some tracks were also steamed at night. Data were then stored in raw format and post-processed using SonarDataEchoview software (Myriax Ltd.) (Higginbottom et al, 2000). All echograms were first scrutinized, the bottom line incorporated, and also background noise was removed according to De Robertis and Higginbottom (2007). Fish abundance was calculated with the 38 kHz frequency as recommended at the PGAAM (ICES 2002), although echograms from 18, 70, 120 and 200 kHz frequencies were used to visually discriminate between fish and other scatter-producing objects such as plankton or bubbles, and to distinguish different fish species according to the frequency response. The 18, 70, 120 and 200 kHz frequencies were used to create a mask allowing a better discrimination between swimbladder fish species and other organisms. The threshold used to scrutinize the echograms was -70 dB. The integration values were expressed as nautical area scattering coefficient (NASC) units or s_A values ($m^2 \text{ nm}^{-2}$) (MacLennan et al., 2002).

1 NASC allocation

A pelagic gear gloria HOD 352 was used to identify the species and size classes responsible for the acoustic energy detected and to provide samples. Haul duration was variable and ultimately depended on the number of fish that enters the net and the conditions where fishing takes place although a minimum duration of 20 minutes was always attempted.

The quality of the hauls for ground-truthing of the acoustic data was classified on account of weather condition, haul performance and the catch composition in numbers and the length distribution of the fish caught as follows (table 2):

Table 2. Ground-truth criteria for fishing stations

	0	1	2	3
Gear performance	Crash	Bad geometry	Bad geometry	God geometry
Fish behaviour		Fish escaping	No escaping	No escaping
Weather conditions	Swell >4 m height Wind >30 knots	Swell: 2-4 m Wind: 30-20 knots	Swell: 1-2m Wind 20-10 knots	Swell <1 m Wind < 10 knots
Fish number	total fish caught <100	Main species >100 Second species <25	Main species > 100 Second species < 50	Main species > 100 Second species > 50
Fish length distribution	No bell shape	Main species bell shape	Main species bell shape Seconds: almost bell shape	Main species bell shape Seconds: bell shape

Hauls considered as the best representation of the fish community for a specific area were used to allocate NASC of each EDSU within this area when no direct allocation was feasible. This process involved the application of the Nakken and Dommasnes (1975, 1977) method for multiple species, but instead of using the mean backscattering cross section, the full length class distribution (1 or 0.5 cm length classes) has been used, as follows:

$$NASC_l = NASC \cdot \left(\frac{\sigma_{l,\rho}}{\sigma_\rho} \right)$$

where $NASC$ is the total backscattering energy to calculate densities by length, $NASC_l$ is the proportion of the total $NASC$ which can be attributed to length group l for a particular fish species. $\sigma_{l,\rho}$ is the backscattering cross-section at length l for a particular species at length l multiplied by the proportion of (ρ_l) of length of this particular species on the overall catch and σ_ρ is the sum of all $\sigma_{l,\rho}$ for all species,

$$\sigma_{l,\rho} = \rho_l * \sigma_l$$

$$\sigma_\rho = \sum_l \sigma_{l,\rho}$$

finally σ_l is backscattering cross-section (m^2) for a fish of length l for a particular species and is computed as follows:

$$\sigma_l = \frac{l \left(\frac{m}{10} \right) * 10 \left(\frac{b_{20}}{10} \right)}{4 * \pi}$$

This is computed from the formula $TS = 20 \log_{L_T} + b_{20}$ (Simmonds and MacLennan, 2005), where L_T is the length class. The b_{20} values for the most important species present in the surveyed area are shown in following table:

Table 3.- b_{20} values from the length target strength relationship of the main fish species assessed in PELACUS survey (WHB is blue whiting; MAC-mackerel; HKE- hake; HOM- horse mackerel; PIL-sardine; JAA-blue jack mackerel (*Trachurus picturatus*); BOG-bogue (*Boops boops*); VMAS-chub mackerel (*Scomber colias*); BOC-board fish (*Capros aper*); and HMM-Mediterranean horse mackerel (*Trachurus mediterraneus*))

Sp	b_{20}	Ref	Observations	Other b_{20}	Ref.
PIL	-72.6	Degnbol et al., 1985	TS for clupeids	-71.2 -70.4 -74.0 -72.5	ICES ,1982 Patti et al., 2000 Hannachi et al., 2005 Georgakarakos et al., 2011
ANE	-72.6	Degnbol et al., 1985	TS for clupeids	-71.2 -76.1 -71.6 -74.8	ICES 1982 Barange et al., 1996 Zhao et al., 2008 Georgakarakos et al., 2011
HKE	-67.5	Foote et al., 1986; Foote, 1987		-68.5 -68.1	Lillo et al., 1996 Henderson, 2005; Henderson and Horne, 2007
BOG	-67.5	Foote et al., 1986	Adapted from gadoids		
BOC	-66.2	Fässler et al., 2013			
MAC	-84.9	Edwards et al., 1984; ICES, 2002		-86.4 -88.0	Misund and Betelstad, 1996 Clay y Castonguay, 1996
HOM	-68.7	Lillo et al., 1996		-68.15 -66.8 -66.5/ 67.0(*)	Gutiérrez and McLennan, 1998 Barange et al. (1996) Georgakarakos et al., 2011
VMA	-68.7	Lillo et al., 1996	Adapted from HOM;l (Sawada, com. pers.)	-70.95	Gutiérrez and McLennan, 1998
WHB	-65.2	Pedersen et al., 2011			

* day and nightrespect.

When possible, direct allocation was done, accounting for the shape of the schools and also the relative frequency response (Korneliussen and Ona, 2003, De Robertis et al, 2010).

Fish schools were extracted using the following settings:

Sv threshold	-60/-70 dB for all frequencies
Minimum total school length	2/20 m
Min. total school height	1/5 m
Min. candidate length	1 m
Min. candidate height	0.5 m
Maximum vertical linking distance	2/5 m
Max. horizontal linking distance	10/25 m
Distance mode	Vessel log
Main frequency for extraction	38/120 kHz

Table 4: Main morphological and backscattering energy characteristics used for schools detection

For all school candidates, several of variables were extracted, among them the NASC (s_A , m^2/nmi^2) together with the proportioned region to cell (ESDU, 1 nmi) NASC and the s_V mean and s_V max and geographic position and time. PRC_NASC values were summed for each ESDU and distances were referenced to a single starting point for each transect. Results for 38 and 120 kHz were compared. Besides, the frequency response for each valid school (i.e. those with length and s_V which allows them be properly measured) was calculated as the ratio $s_{A(f_i)}/s_{A(38)}$, being f_i the s_A values for 18, 70, 120 and 200 kHz.

2 Echointegration estimates

Once backscattering energy is allocated to fish species, the spatial distribution for each species is analysed taking into account both the NASC values and the length frequency distributions (LFD) to provide homogeneous assessment polygons. These are calculated as follows: an empty track determine the along-coast limit of the polygon, whilst three consecutive empty ESDU determine a gap or the across-coast limit. Within each polygon, the LDF is analysed.

LFD were be obtained for all positive hauls for a particular species (either from the total catch or from a representative random sample of 100-200 fish). For the purpose of acoustic assessment, only those LFD which are based on a minimum of 30 individuals will be considered. Differences in probability density functions (PDF) will be tested using Kolmogorov-Smirnov test. PDF distributions without significant differences will be joined, providing a homogeneous PDF strata. Spatial distribution will be then analysed within each stratum and finally mean s_A value and surface (square nautical miles) will be calculated using a GIS based system (Q-gis). These values, together with the length distributions, will be used to calculate the fish abundance in number as described in Nakken and Dommasnes (1975) (see previous section for further details). Estimates for each species will be done on each strata (polygon) using the arithmetic mean of the backscattering energy (NASC, s_A) attributed to each fish species and the surface expressed in square nautical miles using the following formula:

$$\rho_l = \frac{NASC_l}{\sigma_l}$$

$$N_l = \rho_l * A_p$$

where ρ_l is the areal density of fish (numbers per square nautical mile in length group l); the total number for length group l (N_l) within each strata is calculated as the product of ρ_l times the total surface of the strata (A_p) expressed in square nautical miles.

Numbers were converted into biomass using the length weight relationships derived from the fish measured on board. For purposes of comparison, results are given by ICES Sub-Divisions (9aS, CS, CN and N).

3 Centre of Gravity

For each main specie, a centre of gravity (Woillez et al. 2007) was calculated as a weighted average of each sample location (allocated NASC value as weighting factor). Due to the particular topography, instead longitude and latitude, we have used depth and a new variable called "distance from the origin", where the distance (nautical miles) is calculated as $(Lat-37.0)*60$, being Lat the latitude of the middle point of any particular EDSU.

Fishing stations

Fishing stations were used for both NASC allocation and length analysis. Therefore, they will be located on account the results obtained during the acoustic prospection (i.e. opportunistic accounting the echotraces).

A gloria HOD 352 with a vertical opening of about 14 m and around 30 m horizontal one was used. As general rig, 200/400 kg of clump weight were put at each side of the set back (2 m lower wing). Dyneema bridles (wings) have 70 m, but shorten to 50 m in shallower waters. Besides a set of Apollo polyice doors with 3.5 m² and 750 kg weight will be used. Besides, gear performance will be

controlled using a wired Simrad Sonar FS20 net sounder.

Fishing stations were mainly performed during daytime but, exceptionally, some tows were conducted at sunset.

Plankton and hydrological characterisation

Continuous records of SSS, SST and fluorometry are taken using a SeaBird Thermosalinograph coupled with a Turner Fluorometer. Plankton CTD casts down were performed once the acoustic and fishing operations are finished (i.e. from sunset). Four stations are placed over the transects, which are those of the acoustic prospecting but that are extended in order to ensure a minimum distance of 3-6 nmi between stations. The stations are evenly distributed over the surveyed area at a distance of 16-24 nmi.

Zooplankton will be sampled using Bongo net, fitted with 200 and 500 μm nets. The samples will be stored for further analysis at lab.

Top predator observations

Two observers placed at the bridge of the vessel at a height of 16 m above sea level work in turns of two prospecting an area of 180° (each observer covers a field of 90°). Observations are carried out with the naked eye although binoculars are used (7x50) to confirm species identification and determine predator behaviour. Observations are carried out during daylight while the vessel prospects the acoustic transects. Observers record species, number of individuals, behaviour, distance to the vessel and angle to the trackline and observation conditions (wind speed and direction, sea state, visibility, etc.). Observers also record presence, number and type of boats and type, size and number of floating litter. The same methodology is used on the PELGAS surveys and both observer teams shared a common database.

Fish sampling

Catches from fishing trawl hauls were sorted and weighted. All fish species were measured (total length, 1cm classes for all species except clupeids measured at 0.5 cm). When needed, random subsamples of 80-200 specimens were taken. For the main species an additional biological sampling was done for weight, age, sex, maturity stage analysis, complemented by stomach contents analysis (sardine and anchovy); and, sampling for estimation of fecundity adult parameters (sardine). Besides, specific sampling will be done on sardine for pollution and genetic purposes.

1 Catch and length distribution per species

Once sorted the catch, for all species, a length distribution was estimated. If the number of specimens caught was above 100, a random sample was selected. This sample was weighted and the specimens were measured to length class. This was 0.5 for sardine and anchovy and 1 cm for the rest of the species. Catch length distribution was estimated by raising the sample length distribution according to the weighting factor TCW/TSW (total catch weight vs total sampling weight).

2 Weight Length relationship

To all assessed species, a weight length relationship was calculated, either from the results of the biological sampling (see below) or from a specific sampling procedure. In the latter case, a stratified random sampling scheme was used, with the length class (i.e. 0.5 or 1 cm) as stratum.

3 Biological sampling

For main target species caught in each trawl haul (e.g. anchovy and sardine), a biological sampling

was conducted. Data collected were: Length (mm); Weight (g); Sex; Maturity stage; otolith release; fat content; Stomach colour and repletion state. For sardine, the tale will be also collected for further genetic analysis.

RESULTS

The survey was carried out as foreseen. During the first days, NE wind regime was prevalent, which made difficult to perform bongo stations around Galician area; after this episode, compatible with the normal upwelling events in this area, weather was calm and it was only interrupted by an active front with heavy rain during the last weekend. After this front the last 4 days weather was unstable with an increasing strength of the NW wind and swell.

HYDROGRAPHIC CONDITIONS

Accounting the meteorological conditions above described, figure 5 shows the integrated values of surface salinity and temperature.




Figure 5: SST and SSS during IBERAS 0919

As expected, cold surface waters, mainly due to the summer upwelling regime, were located in the northern part (e.g. Aveiro northwards) while the fresher waters were found around Tagus area, due to the front that left heavy rain in this area.

60 plankton station were analysed and the biovolume ($\text{ml}/10\text{m}^3$) from 200 μm and 500 μm , mesh size net determined. The highest abundance were located around the Douro river (figure 6). It should be also highlighted the presence of crustacean, mainly belonging to Euphausiidae. The swarms were identified by fishing stations, as shown in figure 7.




Figure 6: Bongo stations. Left panel (circles encompassed the volume -max 89 ml/m³-) and contours (right) during IBERAS 0919





Figure 7: Echogram at 38 (left), 120 (middle) and 200 kHz (right) of a krill school and its frequency and threshold response (below).

ACOUSTIC

School extraction and total backscattering energy

A total of 6286 echotraces were extracted, accounting for a total NASC (s_A) of 785176 $m^2 nmi^{-2}$. On tracks, NASC values were 430069 $m^2 nmi^{-2}$, which was similar to that recorded in 2018 (476837, a 10 % lower). Figure 1 shows the sum of NASC per track along the surveyed area.






Figure 1. Cumulated NASC values per track

Fish were mainly located in three areas. Between Setúbal canyon and cape Roca; between the mouths of the Mondego and Douro rivers; and within the Rias Baixas. Yet, up to 52% of the total echointegrated energy was found in a single track, round Tocha Beach. The characteristics of these high concentrations will be further analysed.

Most of the schools occurred at 30 m depth, although the maximum cumulated NASC values was found at 22.5 m as shown in figure 2.




Figure 2. Number of schools and their cumulated NASC values per depth strata (5 m)

It seems the main school distribution area was covered as long as only few schools were found in very shallower waters.

Fishing station and echotrace allocation

To perform fishing stations near shore was a challenging task as long as most of the area was occupied by static fishing gears, thus dramatically restricting the available areas to carry out these and increasing the searching time for doing it. In spite this, a total of 26 fishing station were done, accounting a total of 17.4 mt and more than 5.0E+5 specimen as shown in table 5.

Table 5. Summary of the fishing stations (WHB, blue whiting; MAC, mackerel; HKE, hake,; HOM, horse mackerel; PIL, sardine; JAA, bluejack mackerel; BOG, bogue; VMA, chub mackerel; SEAB, seabreams; ANE, anchovy; SNS, longspine snipe fish)

	TOTAL CAP (Kg)	No ind.	No Fishing st	Sample weight (kg	Measured fish	Mean length	%PRES	% Catch_W	% Catch_No
WHB	2	41	1	2	41	19.40	3.85	0.01	0.01
MAC	996	14076	11	17	213	22.48	42.31	5.71	2.64
HKE	2	34	4	2	34	19.00	15.38	0.01	0.01
HOM	3681	70778	22	72	1096	19.65	84.62	21.11	13.28
PIL	1507	32352	18	40	847	17.68	69.23	8.64	6.07
JAA	7	36	1	4	20	0.00	3.85	0.04	0.01
BOG	244	3611	12	28	274	22.43	46.15	1.40	0.68
VMA	4748	73655	18	45	555	22.03	69.23	27.24	13.82
BOC	421	11696	3	10	259	12.44	11.54	2.41	2.20
SEAB	41	128	7	34	109	29.09	26.92	0.24	0.02
ANE	5785	326294	10	7	423	13.71	38.46	33.18	61.24
SNS	2	108	1	1	40	14	3.85	0.01	0.02
Total	17434	532809	26	260	3911				

Horse mackerel was found in 85% of the trawl haul, being also noticeable the presence of sardine and chub mackerel. Anchovy, although was only found in 39% of the stations, accounted for 33% of the total catch in weight (61% in number). That is, had a lower distribution area but was clearly dominant in this. Younger mackerel was also important in the northern part.

Fishing station 26 was not taking into account for allocation purposes. On the other hand, 4 fishing stations were almost monospecific, allowing to identify echotraces.

1 Chub mackerel echotrace identification

Chub mackerel mostly occurred in the southern part. In general, the schools were bigger, isolated, located on the bottom, and very dense. Nevertheless in shallower water, it also occurred in smaller schools. Typical chub mackerel echotrace and its frequency response is shown in figure 3.




Figure 3. Cleaned echogram showing an isolated echotrace attributed to chub mackerel and its characteristic frequency response

11 schools were identified, accounting for a total of 11719 m² nmi⁻². It should be noted that the frequency response together with the schools size and energy descriptors matched with those found during the PELACUS survey.

2 Boar fish echotrace identification

Boar fish was mainly located around the Setúbal canyon. The echotracés, as already seen in the spring surveys in the Spanish area, occurred either close to the bottom, and rising towards middle waters, in isolated schools. On the bottom, it may occur with horse mackerel and sardine.




Figure 4. Cleaned echogram showing an isolated echotrace attributed to boarfish I and its characteristic frequency response

In this case, 10 schools were identified, accounting for a total of 13704 m² nmi⁻².

3 Anchovy echotrace identification

Typically, anchovy occurs in small, continuous echotraces, near bottom, with other fish species. This make difficult to isolate them, and, therefore, rather than direct allocation, fishing stations are used to assign NASC values. In this survey, a big echotrace was detected near Tocha beach. This was a mega-school, recorded in two consecutive tracks, one steamed at sunset and the other in the morning. It has more than 4 nmi long and almost 1 nmi width with a height of about 17 m (i.e. from bottom to sea surface and very thick, as shown in figure 5). The trawl haul performed on this was almost anchovy monospecific with few sardine probably caught in small schools detected before the mega-school (figure 5).




Figure 5. Echogram showing the mega-school at sunset (left) and in the morning (right)

In this case, given that sardine would be also within this school, no direct allocation was done and the energy was split into both species accounting the proportion found at the fishing station

4 Sardine fish echotrace identification

Few echotraces looking like sardine schools in terms of their morphological and energy descriptors were found. The biggest were located in the northern part, within the Murosías.




Figure 6. Echogram showing several echotraces attributed to sardine

12 sardine schools were identified, accounting for a total 36796 m² nmi⁻².

5 Fishing station used for echotrace allocation

From the total of 476837 m² nmi⁻², 75733 were directly allocated to fish species, of which 250346 corresponded to the mega-school, split into fish species on account the result of the fishing station

17, and the rest of the NASC values (150379) were distributed through the species on account the fish proportion found at the fishing stations. Figure 7 shows the spatial distribution of the fishing stations and the proportion for each species estimated using the Nakken and Dommasnes method.

The 9aCS was dominated by chub mackerel while in 9aCN anchovy was predominant and in 9aN horse mackerel which was also important in northern part of 9aCS (near Peniche and Nazaré).

For allocation purposes, the area was split in different strata, on account the *echotypes*. These are areas in which the echotraces were similar and the species proportion found at the fishing station performed on each stratum were also similar. Accordingly, the surveyed area was divided in 11 strata on account echotypes as follows:

1. Near shore community in Algarve, composed by several species, being bogue and chub mackerel dominant (fishing stations 1 and 5)
2. Shelf community in Algarve, mainly composed by boarfish and sardine with a small amount of horse mackerel (fishing station 2)
3. Near shore community in Sado area, with big variety of species but dominated by chub mackerel (fishing stations 4, 5, 6 and 7)
4. Shelf community in Sado, mainly composed by boarfish and sardine with a small amount of horse mackerel (fishing station 8)
5. Near shore community in Lisboa-Caparica area, similar to the Sado community but with bogue as important species. (fishing stations 9 and 10)
6. Community between Ericeira and Nazaré, with sardine and horse mackerel as main species (fishing stations 12 and 13).
7. Community between Nazaré and Figueira da Foz, with anchovy and horse mackerel as main species (fishing stations 11 and 12).
8. Near shore community in Figueira da Foz Porto area with pelagic schools of sardine, horse mackerel and anchovy together with other close to the bottom (fishing stations 11 and 14 and 15). Results from the fishing station 11 were used instead those of number 19 because this was horse mackerel monospecific and thus would likely overestimate the presence of this specie.
9. Shelf community in Figueira da Foz north Aveiro with rather near bottom schools of anchovy and sardine (fishing stations 16, 18 and 20).
10. Shelf community in north Aveiro north Porto with rather near bottom schools of anchovy and sardine, including schools of small sardine (fishing stations 18, 20 and 21).
11. Community in Galician waters, mainly composed by horse mackerel (fishing stations 22, 23, 24 and 25).




Figure 7. Fish proportion accounting the Nakken and Dommaness method (BWH, blue whiting; MAC, mackerel; HAK, hake,; HOM, horse mackerel; PIL, sardine; JAA, bluejack mackerel; BOG, bogue; MAS, chub mackerel; SEAB, seabreams; ANE, anchovy; SNS, longspine snipe fish; PET, small sardine)

Figure 8 shows the amount of energy per echotype as well as the number of nautical miles in which these were found and also the % of energy allocated to each main species. On overall, 29% of the total energy was allocated to horse mackerel, while for chub mackerel this percentage was a 33% and for anchovy a 24%. For sardine the percentage was a 7%. Echotype 3 (the one characterised by the fishing stations 4, 5, 6 and 7), accounted for more than the 30% of the total energy being the bulk of this allocated to chub mackerel; in the same way in north Portugal, main echotypes were mainly characterised by anchovy while in the central part and in Galicia, the major contribution was due to horse mackerel.




Figure 8. Left, in solid blue bars, NASC allocated (%) to each echotype and in dashed bars, the number of nautical miles for each echotype. Right, fish proportion accounting the Nakken and Dommaness method to each echotype (ANE, anchovy; BOG, bogue; VMA, chub mackerel; PIL, sardine; and HOM horse)

Figure 9 shows the spatial distribution of each echotype strata.




Figure 9. Spatial distribution of each echotype strata. Numbers reflect the same numbering of the echotype description. Number 12 deals with the area of the mega-school

Acoustic assessment

Table 6 shows the total energy attributed to the main species as well as the center of gravity, using as coordinates the distance from the origin, located at 37°N, and depth. As expected, most of the integrated backscattering energy was attributed to anchovy, being also important those for horse mackerel, chub mackerel and sardine. 76% of the sardine NASC was directly allocated. It was also important the direct allocation done for boar fish (88%) while for the rest of the species this % was lower than 25%.

	MAC	HOM	PIL	JAA	BOG	VMA	BOC	ANE	SNS	PET
NASC	278	57152	48423	45	6335	61378	15493	285688	18	302
Depth	27.07	31.54	20.04	69.04	31.28	30.53	76.62	17.72	69.04	26.73
s.d.	3.24	54.17	23.51	1.13	18.35	48.32	37.14	31.36	0.72	1.10
ic	0.52	8.66	3.76	0.18	2.93	7.73	5.94	5.01	0.12	0.18
Dist	225.46	250.74	234.98	78.52	114.40	82.88	87.14	203.40	78.52	210.66
s.d.	11.43	298.61	335.28	0.45	94.84	53.19	58.48	62.58	0.29	5.55
ic	1.83	47.75	53.61	0.07	15.16	8.50	9.35	10.01	0.05	0.89

Table 6. Total NASC allocated to the main pelagic species together with the location of the coordinates of the centre of gravity (MAC, mackerel; HOM, horse mackerel; PIL, sardine; JAA, blue jack mackerel; BOG, bogue; VMA, chub mackerel; BOC, boarfish, ANE, anchovy; SNS, longspine snipefish, PET, small sardine)





Figure 10. Center of gravity and cumulated NASC for the most important pelagic species (ANE, anchovy-green-; PIL, sardine -blue-; HOM, h. mackerel -yellow-; MAC, mackerel -red-; VMA, C. mackerel -orange-; BOC, boarfish -black-; and BOG, bogue -magenta-)

Figure 10 shows the spatial distribution of the center of gravity as well as the cumulated NASC along distance from the origin. Chub mackerel and bogue were mainly located in 9aCS, round Setúbal area, together with boarfish which has its center of gravity in the outer area (deeper waters). Those of sardine, anchovy, mackerel and horse mackerel were found closer each other, and located in 9aCN, between Mondego and Douro rivers, where it seems their main recruitment area is placed.

1 Sardine assessment

Accounting the length distributions obtained at the fishing station and the NASC spatial distribution, sardine was divided in 6 strata, 4 in 9aCS and a single stratum in both 9aCN and 9aN. In the Setúbal strata, mean length was lower than those of the surrounding areas (north and south) and also length distribution was significantly different. Besides there was a discontinuity between Ericeira and Lisboa strata (with only 2 positive miles with NASC values lower than $1 \text{ m}^2 \text{ nmi}^{-2}$); both the difference in mean length/length distribution and the discontinuities in the fish distribution explained the strata. In 9aCN all fishing station gave a lower mean length than those estimated in 9aN and in Ericeira area.

Table 7 summarises the sardine assessment. A total of 35701 tonnes, corresponding to 819 million fish were estimated. The bulk of the distribution was found in 9aCN ($16.6 \cdot 10^3$ tonnes).

ICES-Div	Region	SURVEY: IBERAS 1118 SARDINE			Fishing st.	PDF	No (million f ish)	Biomass (tonnes)	Density (Tn/nmi-2)
		No	Mean	Surface					
9aCS	Sines	44	132.04	181	P02-P04-P07-P09-P10-P11	S01	86	6035	33
	Setúbal	37	25.03	105	P07-P08	S02	14	515	5
	Lisboa	42	35.93	138	P02-P04-P07-P09-P10-P11	S01	18	1249	9
	Ericeira	52	21.53	394	P02-P04-P07-P09-P10-P11	S01	31	2144	5
	Total	175	53.51	818			149	9944	12
9aCN	9aCN	178	110.13	856	P14-15-P16-P17-P18-P20	S03	581	16604	19
	Total	178	110.13	856			581	16604	19
9aN	Rias Baixas	68	290.37	112	P22	S04	88	9153	82
	Total	68	290	112			88	9153	82
	Total Portugal	353	82	818			730	26548	32
	Total Spain	68	290	112			88	9153	82
	TOTAL 9a	421	115.71	930			819	35701	38

Table 7. Summary of the sardine assessment, with the name of the strata, number of positive nmi, mean NASC value ($m^2 nmi^{-2}$), surface (nmi^2), fishing station used for the estimation and number and biomass estimated

Length distribution were quite complementary among areas, with the younger fish located in 9aCN, the older in 9aN and the intermediate lengths in 9aCS, as shown in figure 11.

Figure 12 shows the spatial distribution accounting the NASC values. Main distribution area seems to be located around Figueira da Foz, being similar that observed for sardine. Yet, two second areas were located around Setúbal and within the RíasBaixas (9aN) in Spain, in which anchovy was almost absent. Between Aveiro and Porto, the distribution extended off shallower waters, and the outer limit of the sardine distribution was not reached.




Figure 11. Sardine estimated abundance and biomass per length class in 9aCS (above left), 9aCN (above right), 9aN (below left) and for the total area (below right)




Figure 12. Sardine spatial distribution in IBERAS 1118. Dots represent the NASC values attributed to sardine and the polygons the strata together with the relative density

On the other hand, sardine adult distribution matched quite well with the egg distribution derived from the CUFES samples, as shown in figure 13.




Figure 13: Sardine egg spatial distribution in IBERAS 1118 obtained from CUFES samples

Table 8a-e is shown the sardine assessment by length group and age classes per ICES Sub-Division and by country. All recruit (age group 0) were found in the Portuguese area, accounting for the 73.44% of the total fish estimates in this country. Moreover, the bulk of them were recorded in 9aCN where juveniles accounted up to 90% of the total estimates in number. O group mean length was 14.83 in 9aCN, similar to that found in 9aCS (14.80), although in 9aCN a small second mode was also recorded at 10.5 cm.

Age group 1 was almost negligible, confirming, thus, the very weak strength of this cohort.

In 9aN (Spain), only older fish were observed, without younger fish. It should be noticed that in this area the lack of time together with the bad weather conditions found at the end of the survey made difficult to obtain extra trawl hauls samples in order to verify the true length structure.

SURVEY: IBERAS 0318. Sardine

BIOMASS (tonnes). ZONE: 9aCS

Length	AGE GROUPS										Total	No fish (thousands)	
	0	1	2	3	4	5	6	7	8	9			
10													
10.5													
11													
11.5													
12													
12.5													
13	11											11.09	644
13.5	25											25.12	1288
14	78											77.92	3542
14.5	64											63.66	2576
15	36											35.61	1288
15.5	8											7.62	247
16	18											17.83	520
16.5	43											43.32	1142
17	11											11.45	273
17.5			69									69.17	1500
18		38	229									267.27	5276
18.5		33	962	33								1028.07	18528
19			1638	157								1795.22	29606
19.5		35	1672	279	70	35						2090.48	31620
20			1052	137								1189.46	16537
20.5			520	189	47							755.93	9680
21			217	173	87	43	87					607.09	7174
21.5			66	197		131	66					458.76	5012
22					70	139	209		70			486.61	4925
22.5					138	69	207	69				484.07	4545
23					85	85		85				253.85	2215
23.5					82		82					164.06	1332
24													
24.5													
25													
25.5													
26													
26.5													
27													
Biomass (mt)	294	106	6424	1166	578	502	650	154	70	0		9943.66	149472
%	2.95	1.07	64.61	11.72	5.81	5.05	6.54	1.55	0.70				
M. weight	25.01	56.29	63.74	72.59	93.55	94.66	100.19	110.76	98.81			66.08	
No Fish (thousands)	11522	1878	100221	15937	6091	5266	6466	1388	704	0		149472	
%	7.71	1.26	67.05	10.66	4.07	3.52	4.33	0.93	0.47				
M. length	14.80	18.83	19.54	20.31	21.89	21.97	22.34	23.02	22.25			19.59	
s.d.	1.00	0.61	0.74	0.88	1.33	0.94	0.70	0.25				1.84	

Table 8a: Sardine assessment in 9aCS

SURVEY: IBERAS 0318. Sardine

BIOMASS (tonnes). ZONE: 9aCN

Length	AGE GROUPS										Total	No f ish (thousands)
	0	1	2	3	4	5	6	7	8	9		
10	43										43.12	5715
10.5	167										167.07	18983
11	71										71.01	6967
11.5												
12	26										25.98	1936
12.5	147										147.23	9641
13	434										433.83	25091
13.5	846										845.53	43388
14	1195	54									1249.48	57130
14.5	2211										2210.74	90428
15	3023										3022.79	111023
15.5	2041										2040.90	67543
16	1523										1522.81	45558
16.5	777	283									1059.50	28742
17	296	118									414.08	10215
17.5	415	277	46								737.85	16598
18	299	100	150								548.40	11277
18.5		93	155								247.89	4671
19			75								74.62	1292
19.5		29	146	146							321.76	5127
20			300	263	75						638.20	9380
20.5		47	47	236	47						377.30	5125
21			119	79	40						238.46	3000
21.5				55	111						165.90	1936
22												
22.5												
23												
23.5												
24												
24.5												
25												
25.5												
26												
26.5												
27												
Biomass (mt)	13513	1001	1038	780	273	0	0	0	0	0	16604.46	580767
%	81.38	6.03	6.25	4.70	1.64							
M. weight	24.89	40.51	59.74	70.43	76.87						27.75	
No Fish (thousands)	524781	24200	17209	11041	3535	0	0	0	0	0	580767	
%	90.36	4.17	2.96	1.90	0.61							
M. length	14.83	17.25	19.45	20.47	21.03						15.21	
s.d.	1.48	1.33	1.02	0.55	0.63						1.91	

Table 8b: Sardine assessment in 9aCN

SURVEY: IBERAS 0318. Sardine

BIOMASS (tonnes). ZONE: 9aC

Length	AGE GROUPS									Total	No fish (thousands)	
	0	1	2	3	4	5	6	7	8			9
10												
10.5												
11												
11.5												
12												
12.5												
13												
13.5												
14												
14.5												
15												
15.5												
16												
16.5												
17												
17.5												
18												
18.5												
19												
19.5												
20												
20.5												
21												
21.5				665	1330						1995.26	23282
22												
22.5					461	461	461			461	1845.61	18626
23								2475			2474.82	23282
23.5					530						530.17	4656
24				425		425		425		425	1701.22	13969
24.5				303		303					605.71	4656
25												
25.5												
26												
26.5												
27												
Biomass (mt)	0	0	0	1393	2322	1190	461	2900	0	887	9152.79	88472
%				15.22	25.37	13.00	5.04	31.69		9.69		
M. weight				101.32	93.07	113.00	99.09	108.23		108.42	103.02	
No Fish (thousands)	0	0	0	13581	24834	10477	4656	26774	0	8149	88472	
%				15.35	28.07	11.84	5.26	30.26		9.21		
M. length				22.91	22.31	23.69	22.75	23.38		23.39	23.01	
s.d.				1.35	0.79	0.86		0.34		0.74	0.94	

Table 8c: Sardine assessment in 9aN (Spain)

SURVEY: PELACUS 0318. Sardine

BIOMASS (tonnes). ZONE: 9aCS-CN

Length	AGE GROUPS										Total	No fish (thousands)
	0	1	2	3	4	5	6	7	8	9		
10	43										43.12	5715
10.5	167										167.07	18983
11	71										71.01	6967
11.5												
12	26										25.98	1936
12.5	147										147.23	9641
13	445										444.91	25735
13.5	871										870.65	44676
14	1273	54									1327.40	60673
14.5	2274										2274.40	93005
15	3058										3058.40	112312
15.5	2049										2048.52	67790
16	1541										1540.64	46079
16.5	820	283									1102.82	29884
17	307	118									425.53	10488
17.5	415	277	115								807.03	18097
18	299	138	379								815.66	16554
18.5		126	1117	33							1275.95	23199
19			1712	157							1869.84	30898
19.5		64	1819	425	70	35					2412.24	36747
20			1353	400	75						1827.66	25917
20.5		47	567	425	94						1133.23	14805
21			336	253	126	43	87				845.55	10174
21.5			66	252	111	131	66				624.66	6948
22					70	139	209		70		486.61	4925
22.5					138	69	207	69			484.07	4545
23					85	85		85			253.85	2215
23.5						82		82			164.06	1332
24												
24.5												
25												
25.5												
26												
26.5												
27												
Biomass (mt)	13807	1107	7463	1945	851	502	650	154	70	0	26548.12	730239
%	52.01	4.17	28.11	7.33	3.20	1.89	2.45	0.58	0.26			
M. weight	25.91	41.26	58.38	66.20	78.44	82.73	86.95	94.95	85.91		35.00	
No Fish (thousands)	536303	26079	117431	26978	9625	5266	6466	1388	704	0	730239	
%	73.44	3.57	16.08	3.69	1.32	0.72	0.89	0.19	0.10			
M. length	14.83	17.36	19.52	20.37	21.58	21.97	22.34	23.02	22.25		16.11	
s.d.	1.47	1.35	0.79	0.77	1.20	0.94	0.70	0.25			2.59	

Table 8d: Sardine assessment in Portugal (9aCS+9aCN)

Length	AGE GROUPS									Total	No fish (thousands)	
	0	1	2	3	4	5	6	7	8			9
10	43										43.12	5715
10.5	167										167.07	18983
11	71										71.01	6967
11.5												
12	26										25.98	1936
12.5	147										147.23	9641
13	445										444.91	25735
13.5	871										870.65	44676
14	1273	54									1327.40	60673
14.5	2274										2274.40	93005
15	3058										3058.40	112312
15.5	2049										2048.52	67790
16	1541										1540.64	46079
16.5	820	283									1102.82	29884
17	307	118									425.53	10488
17.5	415	277	115								807.03	18097
18	299	138	379								815.66	16554
18.5		126	1117	33							1275.95	23199
19			1712	157							1869.84	30898
19.5		64	1819	425	70	35					2412.24	36747
20			1353	400	75						1827.66	25917
20.5		47	567	425	94						1133.23	14805
21			336	253	126	43	87				845.55	10174
21.5			66	917	1441	131	66				2619.92	30230
22					70	139	209		70		486.61	4925
22.5					600	531	669	69		461	2329.68	23171
23					85	85		2559			2728.66	25497
23.5					612		82				694.24	5989
24				851		851					1701.22	13969
24.5				303		303					605.71	4656
25												
25.5												
26												
26.5												
27												
Biomass (mt)	13807	1107	7463	3764	3172	2117	1112	2629	70	461	35701	818711
%	38.67	3.10	20.90	10.54	8.89	5.93	3.11	7.36	0.19	1.29		
M. weight	25.91	41.26	58.38	77.21	84.28	98.73	88.94	97.66	85.91	91.74	41.43	
No Fish (thousands)	536303	26079	117431	44052	34460	19236	11122	24670	704	4656	818711	
%	65.51	3.19	14.34	5.38	4.21	2.35	1.36	3.01	0.09	0.57		
M. length	14.83	17.36	19.52	21.46	22.11	23.32	22.51	23.24	22.25	22.75	16.86	
s.d.	1.47	1.35	0.79	1.71	0.98	1.18	0.00	0.00	0.00	0.00	3.27	

Table 8e: Sardine assessment in the whole area (9aCS+9aCN+9aN)

Figure 14 shows the sardine assessment by age group and ICES sub-division, in Portugal and for the whole area. There were remarkable difference in age composition among sub-divisions. While in southern Portuguese Atlantic waters (9aCS) age 2 was clearly dominant, in the northern Portuguese region the bulk of the population in the surveyed area was composed by younger fish (e.g. young of the year), while in the Spanish area, only individuals older than 2 were found.




Figure 14: Sardine assessment (abundance and biomass by age group) by ICES division

2 Anchovy assessment

For anchovy, the 9aCN Sub-Division was divided into 5 strata. 3 of them accounting the differences in length distribution and mean length and the other two to restrict the influence of the mega-school to the adjacent areas; besides, another strata was placed in the northern part of 9a CN because this was mainly steamed during night hours.

Mainly due to the mega-school found north Figueira da Foz, the total biomass estimated was 182 thousand tonnes as shown in table 9. Almost no anchovy was found in 9aN, nor in 9aCS. Main mode was found at 14.5 cm, with a secondary one at 16.5 cm, as shown in figure 15.

SURVEY: IBERAS1118 ANCHOVY

Zone	Area	No	Mean	Area	Fishing st.	PDF	No (million fish)	Biomass (tonnes)
9aCS	Lisboa	39	0.02	134	P14	S01	0	0
	Total	39	0.02	134			0	0
9aCN	Figueira	54	23.14	224	P14	S01	34	670
	Mira	3	83057.20	15	P16-P17-P18	S02	7568	161674
	Aveiro	81	308.13	341	P16-P17-P18	S02	654	13979
	Porto	69	148.85	357	P20-P21	S03	557	5047
	Viana	9	2.15	72	P20-P21	S03	23	206
	Total	216	1322.55	1009			8836	181576
9aN	Rbaixas	68	0.04	112	P20-P21	S03	0	0
	Total	68	0.04	112			0	0
	Portugal	255	1120	1142			8836	181577
	Spain	68	0	112			0	0
	TOTAL	323	884.44	1254			8836	181577

Table 9. Summary of the anchovy assessment, with the name of the strata, number of positive nmi, mean NASC value ($m^2 nmi^{-2}$), surface (nmi^2), fishing station used for the estimation and number and biomass estimated




Figure 15. Anchovy estimated abundance and biomass per length class in 9aCS (above left), 9aCN (above right), 9aN (below left) and for the total area (below right), note that scales are different

Regarding ages, it should be noted that although the high abundance compared to that estimated for sardine, only 5.9 kt belonged to age group 0 (recruits), accounted age group 1 for a 84% of the total biomass. Comparing to sardine where recruits represented up 66% of the total abundance and 39% in biomass, juvenile anchovy only accounted for the 7% of the abundance (3% of the total biomass). Given these figures, it seems 2018 recruitment would be very low as compared to that of 2017. Figure 16 and table 10 shows the anchovy assessment by age group.

Length	AGE GROUPS				Total	No f ish (thousands)	Length	AGE GROUPS				Total	No f ish (thousands)	
	0	1	2	3				0	1	2	3			
5							5							
5.5							5.5							
6							6							
6.5							6.5							
7							7							
7.5							7.5							
8							8							
8.5							8.5							
9							9							
9.5							9.5	30				30.41	5916	
10							10	252				251.82	41414	
10.5							10.5	676				675.51	94660	
11							11	924	257			1180.54	141990	
11.5	0	0			0.00	0	11.5	807	224			1030.78	107119	
12	0	0			0.00	0	12	1689	1014			2702.95	244179	
12.5	0	0			0.01	1	12.5	374	2616			2989.76	236108	
13	0	0	0		0.01	1	13	392	6274	392		7058.64	489831	
13.5	0	0			0.03	2	13.5	751	8256			9006.12	551815	
14		0			0.05	3	14		19080			19080.03	1036805	
14.5		0	0		0.06	3	14.5		26862	3837		30699.50	1485626	
15		0	0		0.09	4	15		25135	2900		28035.36	1212894	
15.5		0	0		0.04	2	15.5		15344	8631		23975.68	930671	
16		0	0		0.03	1	16		20927	1163		22089.87	771967	
16.5		0	0		0.02	1	16.5		23696	4182		27877.99	879893	
17		0			0.01	0	17		7691	0		7690.55	219882	
17.5		0	0		0.01	0	17.5		7554	3022		10576.18	274696	
18		0	0		0.01	0	18		3107	1553		4660.39	110255	
18.5							18.5							
19							19							
19.5							19.5							
20							20							
Biomass (mt)	0	0	0	0	0.40	18	Biomass (mt)	5894	168037	25681	0	199612.10	8835724	
%	2.28	85.82	11.90				%	2.95	84.18	12.87				
M. weight	11.80	21.23	25.38		21.26		M. weight	9.69	22.41	26.25		21.98		
No Fish (thousands)	1	16	2		18		No Fish (thousands)	592269	7282175	961279	0	8835724		
%	4.15	85.83	10.02				%	6.70	82.42	10.88				
M. length	12.48	14.87	15.68		14.85		M. length	11.77	15.11	15.84		14.97		
s.d.	0.80	1.21	1.56		1.30		s.d.	0.97	1.30	1.05		1.54		

Table 10a: Anchovy assessment in 9aCS and 9aCN

Length	AGE GROUPS				Total	No fish ('000)	Length	AGE GROUPS				Total	No fish ('000)
	0	1	2	3				0	1	2	3		
5	0	0	0	0			5						
5.5	0	0	0	0			5.5						
6	0	0	0	0			6						
6.5	0	0	0	0			6.5						
7	0	0	0	0			7						
7.5	0	0	0	0			7.5						
8	0	0	0	0			8						
8.5	0	0	0	0			8.5						
9	0	0	0	0			9						
9.5	30	0	0	0	30.41	5916	9.5	0				0.00	0
10	252	0	0	0	251.82	41414	10	0				0.02	3
10.5	676	0	0	0	675.51	94660	10.5	0				0.05	7
11	924	257	0	0	1180.54	141990	11	0	0			0.09	11
11.5	807	224	0	0	1030.79	107120	11.5	0	0			0.08	8
12	1689	1014	0	0	2702.95	244179	12	0	0			0.11	10
12.5	374	2616	0	0	2989.77	236109	12.5	0	0			0.02	1
13	392	6274	392	0	7058.66	489832	13						
13.5	751	8256	0	0	9006.14	551817	13.5	0	0			0.01	0
14	0	19080	0	0	19080.08	1036808	14						
14.5	0	26862	3837	0	30699.57	1485629	14.5		0	0		0.02	1
15	0	25135	2900	0	28035.46	1212898	15		0	0		0.01	0
15.5	0	15344	8631	0	23975.73	930673	15.5						
16	0	20927	1163	0	22089.90	771968	16		0	0		0.01	0
16.5	0	23696	4182	0	27878.01	879894	16.5		0	0		0.01	0
17	0	7691	0	0	7690.57	219883	17						
17.5	0	7554	3022	0	10576.19	274697	17.5						
18	0	3107	1553	0	4660.41	110255	18						
18.5	0	0	0	0			18.5						
19	0	0	0	0			19						
19.5	0	0	0	0			19.5						
20	0	0	0	0			20						
Biomass (mt)	5894	168038	25681	0	199612.49	8835742	Biomass (mt)	0	0	0	0	0.44	44
%	2.95	84.18	12.87				%	63.91	34.66	1.43			
M. weight	9.69	22.41	26.25		21.98		M. weight	8.51	12.32	24.54		9.60	
No Fish ('000)	592270	7282191	961281	0	8835742		No fish ('000)	32	12	0	0	44	
%	6.70	82.42	10.88				%	73.17	26.27	0.57			
M. length	11.77	15.11	15.84		14.97		M. length	11.33	12.65	15.52		11.70	
s.d.	0.97	1.30	1.05	0.00	1.54		s.d.	0.68	1.55	1.11		1.16	

Table 10b: Anchovy assessment in Portugal and Spain

Length	AGE GROUPS				Total	No fish (thousands)
	0	1	2	3		
5	0	0	0	0		
5.5	0	0	0	0		
6	0	0	0	0		
6.5	0	0	0	0		
7	0	0	0	0		
7.5	0	0	0	0		
8	0	0	0	0		
8.5	0	0	0	0		
9	0	0	0	0		
9.5	30	0	0	0	30.41	5917
10	252	0	0	0	251.84	41417
10.5	676	0	0	0	675.56	94667
11	924	257	0	0	1180.63	142001
11.5	807	224	0	0	1030.86	107128
12	1689	1014	0	0	2703.06	244190
12.5	374	2616	0	0	2989.78	236110
13	392	6274	392	0	7058.66	489832
13.5	751	8256	0	0	9006.15	551817
14	0	19080	0	0	19080.08	1036808
14.5	0	26862	3837	0	30699.59	1485630
15	0	25135	2900	0	28035.47	1212899
15.5	0	15344	8631	0	23975.73	930673
16	0	20927	1163	0	22089.91	771968
16.5	0	23696	4182	0	27878.03	879894
17	0	7691	0	0	7690.57	219883
17.5	0	7554	3022	0	10576.19	274697
18	0	3107	1553	0	4660.41	110255
18.5	0	0	0	0		
19	0	0	0	0		
19.5	0	0	0	0		
20	0	0	0	0		
Biomass (mt)	5894	168038	25681	0	199612.93	8835786
%	2.95	84.18	12.87			
M. weight	9.69	22.41	26.25		21.98	
No Fish (thousands)	592302	7282203	961282		8835786	
%	6.70	82.42	10.88			
M. length	11.77	15.11	15.84		14.97	
s.d.	0.97	1.30	1.05		1.54	

Table 10c: Anchovy assessment in the whole area (9aCS+9aCN+9aN)




Figure 16: Anchovy assessment (abundance and biomass by age group) by ICES division

Finally, figure 17 shows the spatial distribution.




Figure 17. Anchovy spatial distribution in IBERAS 1118. Dots represent the NASC values attributed to anchovy and the polygons the strata together with the relative density

3 Horse mackerel assessment

Horse mackerel was well distributed all around the surveyed area but in in the southern part (Algarve). Length distribution were similar around the surveyed area, noticing the presence of very small juveniles in hauls, specially in 9aCS and 9aN. 9aCS was divided into three strata due to a discontinuities in the spatial distribution, although length distribution were similar throughout the Sub-Division. On the other hand, in 9aCN an extra strata was placed in the northern part as this was mainly steamed at night. Table 11 summarises the assessment. A total of 22 thousand tonnes were estimated, corresponding to 341 million fish.

Most of the younger fish occurred in 9aCN, while only few juveniles were recorded in 9aCS. In 9aN, a mode peaked in 12 cm, not seen in 9aCN was also recorded although most of the fish were bigger than 20 cm. (figure 18)

Figure 19 shows its spatial distribution and the relative density. North Figueira da Foz and within the Rias Baixas, the density was the highest.

SURVEY: IBERAS 1118 HORSE MACKEREL

Zone	Area	No	Mean	Surface	Fishing st.	PDF	No (million fish)	Biomass (tonnes)
9aCS	Alentejo	10	4.22	91.27	-P02-P03-P04-P05-P06-P08-P09-	ST01	0	49
	Lisboa-Sines	126	30.04	479.79	-P02-P03-P04-P05-P06-P08-P09-	ST01	16	1841
	Peniche	52	38.09	403.54	-P02-P03-P04-P05-P06-P08-P09-	ST01	17	1964
	Total	188	31	974.60			33.88	3853.84
9aCN	Aveiro	197	125.38	899.54	P11-P12-P15-P16-P19-P21	ST02	191	11060
	Viana	23	50.51	183.13	P11-P12-P15-P16-P19-P21	ST02	16	907
	Total	220	118	1082.7			206.85	11967.59
9aN	Rbaix as	118	215.93	293.68	P22-P23-P24-P25	ST03	100	6511
	Total	118	216	294			99.81	6511.01
Total Portugal		408	78	2057			241	15821
Total Spain		118	216	294			100	6511
Total 9a		526	109	2351			341	22332

Table 11. Summary of the horse mackerel assessment, with the name of the strata, number of positive nmi, mean NASC value ($m^2 nmi^{-2}$), surface (nmi^2), fishing station used for the estimation and number and biomass estimated




Figure 18. Horse mackerel estimated abundance and biomass per length class in 9aCS (above left), 9aCN (above right), 9aN (below left) and for the total area (below right)




Figure 19. Horse mackerel spatial distribution in IBERAS 1118. Dots represent the NASC values attributed to horse mackerel and the polygons the strata together with the relative density

4 Chub mackerel assessment

Table 12 summarises the chub mackerel assessment. Of 25 thousand tonnes, 99% were found in 9aCS. The distribution was bimodal, with the main mode at 19 cm and the second one at 23 cm (figure 20). Figure 21 shows the spatial distribution, with the bulk of the distribution being located close to the Sado estuary.

SURVEY: IBERAS 1118 CHUB MACKEREL									
Zone	Area	No	Mean	Surface	Fishing st.	PDF	No (million fish)	Biomass (tonnes)	
9aCS	Alentejo	10	46.08	91.27	-P04-P05-P06-P07-P08-P09-P10-I	ST01	5	438	
	Lisboa-Sines	126	473.43	479.79		ST01	274	23674	
	Peniche	52	18.19	403.54		ST01	9	765	
	Total	188	325	974.60				288.08	24877.42
9aCN	Aveiro	197	1.37	899.54	P09-P10-P12	ST02	1	130	
	Viana	23	0.40	183.13	P05-P21	ST03	0	10	
	Total	220	1	1082.7			1.48	140.34	
9aN	Rbaixas	118	0.10	293.68	P05-P21	ST03	0	4	
	Total	118	0	294			0.02	3.90	
Total Portugal		408	150	2057			290	25018	
Total Spain		118	0	294			0	4	
Total 9a		526	117	2351			290	25022	

Table 12. Summary of the chub mackerel assessment, with the name of the strata, number of positive nmi, mean NASC value ($m^2 nmi^{-2}$), surface (nmi^2), fishing station used for the estimation and number and biomass estimated




Figure 20. Chub mackerel estimated abundance and biomass per length class in 9aCS (above left), 9aCN (above right), 9aN (below left) and for the total area (below right), note that scales are different




Figure 21. Chub mackerel spatial distribution in IBERAS 1118. Dots represent the NASC values attributed to chub mackerel and the polygons the strata together with the relative density

DISCUSSION AND CONCLUSIONS

IBERAS1118 is the first attempt for surveying the main potential area of the recruitment of the iberatlantic stock of sardine using the new EK80 echosounder on board of the Spanish Research vessel Ramón Margalef. The survey was carried out by an integrated team composed by scientists and technician of both IPMA (Portugal) and IEO (Spain) research centres.

In general, weather conditions made difficult to undertake all the foreseen activities. In spite of this, the expected acoustic coverage was performed together with the fishing stations. However, some concerns could arise due to the impossibility for surveying the coastal waters for safety reasons due to the height of the swells. Nevertheless, on account figure 2, the school distribution along depth showed that the bulk of them were located at around 30 m, within the depth range of the surveyed area. Besides, the bad weather conditions, with heavy rain left the coastal waters less saltier and fish use to avoid these kind of waters. In fact, in 9aN, the bulk of the 0 group seemed to be located really far from the coastal areas. Even more, in 2017, mean depth of the sardine abundance was estimated at 25.90 m and 17.09 for anchovy. This year was 20.04 and 17.72, which means, mean depth remained stable for anchovy and more in shallower waters for sardine but still inside the surveyed area. On account these results, it seems only few fish would be located near shore, thus, outside the vessel operative range, so the achieved results are comparable to those of the 2017 survey (JUVESAR).

Comparing both survey, it was a clear change in the distribution pattern of sardine, as shown in figure 22.




Figure 22. Sardine spatial distribution in JUVESAR 17 (left) and IBERAS 1118 (right). Dots represent the NASC values attributed to sardine

In 2017, sardine showed a very patchy and dense distribution with only few hot spots (six in total), while in 2018, although some dense patches, the distributions was wider, showing a continuity among transects.

For anchovy this feature was even more clear, with a clear expansion of the distribution area. As

shown in figure 23.




Figure 23. Anchovy spatial distribution in JUVESAR 17 (left) and IBERAS 1118 (right). Dots represent the NASC values attributed to anchovy

Probably, the number of hotspots with high fish concentration is a feature during winter time, with concentrating in smaller areas, and spreading its distribution area towards wider areas in spring time, when food availability increases. In northern areas, herring and mackerel among other species exhibit this kind of behaviour, with most of the fish concentrated in few but big schools (Huse and Ona 1996). During this survey anchovy, and in lesser amount chub mackerel, occurred in big and almost isolate schools. These big schools increase the uncertainty by decreasing the precision as long as they have an important impact in both overall mean abundance and its variance estimates. In order to avoid this, it should be better to perform the survey earlier, (i.e. beginning fall, end summer), when fish would occur in a wider area. Besides, if the survey is performed in September instead November, the survey will take place at the same time of the JUVENA survey, thus giving a synoptic coverage of the Iberian Peninsula. In any case, placing the intertransect distance at 4 nmi apart, sounds reasonable given both the aggregation and the spatial distribution patterns.

During IBERAS survey, although the estimated biomass of anchovy was much higher than that of the sardine (round 200 kt and 35 kt respectively), it should be noted that the biomass of the young fish (recruitment) was higher in sardine (round 14 kt) than that assessed for anchovy (only 6 kt). Although the anchovy spawning season takes place later in the year (e.g. March -June), in the Bay of Biscay pre-recruits are found even in September in open waters and accessible to the survey vessel, so the recruitment would be also accessible to the surveyed area unless, as in the northern part, it is unavailable because is located off-survey track.

Comparing with previous juvenile target surveys in Portugal (JUVESAR time series), the number of sardine juveniles in 2018 was higher than those estimated in 2017 (525 million fish in 2018 and 472 million fish in 2017, see table 13), although the biomass was higher in 2017 (1 kt more). In any case, these figures are the lowest of the time series 2013-18. Curiously, the assessed anchovy biomass in

2018 is the highest of the time series but the strength of the recruitment is the weakest of the time series.

The spring surveys PELAGO and PELACUS will confirm in any case the strength of both sardine and anchovy recruitment.

Survey	Sardine				Anchovy			
	Total (millions)	Total (kt)	Recruits (millions)	Recruits (kt)	Total (millions)	Total (kt)	Recruits (millions)	Recruits (kt)
<i>Série Juvesar</i>								
JUVESAR13	2 093	22	1 986	19,4	-	-	-	-
JUVESAR15	2 831	45	2 822	44,4	3 870	30,0	3 835	29,0
JUVESAR16	1 264	57	899	28,6	2 836	14,4	2 835	14,4
JUVESAR17*	2 077	120	472	14,6	2 145	38,0	570	4,7
IBERAS/JUV18 +	629,3	20	525	13,5	8 836	181,6	391	2,9
<i>Extended area</i>								
JUVESAR17**	2 292	132	472	14,6	2 145	38,0	570	4,7
IBERAS/JUV18 ++	643,8	20,5	535,3	13,8	8 836	181,6	391	2,9
IBERAS/JUV18+++	818,7	35,7	536,3	13,8	8 836	181,6	391	2,9

Table 13 – Sardine and anchovy abundance and biomass estimates in the November time series 2013-18 (*only in the area between Espichel Cape and northern boundary; + area between Caminha and Espichel Cape for interannual comparison purposes; ** from Póvoa to Melides; ++ from Caminha to Sagres; +++ whole surveyed area (from Galiza to Sagres). Recruits mean age 0 for sardine and L < 12 cm for anchovy.

CONSULTED BIBLIOGRAPHY

- Barange, M., Hampton, I. And Soule, M. 1996 Empirical determination of in situ target strength of three loosely aggregated pelagic fish species. *ICES Journal of Marine Science*, 53: 225-232.
- Barange, M., and Hampton, I. 1997. Spatial structure of co-occurring anchovy and sardine populations from acoustic data: implications for the survey design. *Fisheries Oceanography*, 6(2): 94-108.
- Boyra, G., Martínez, U., Cotano, U., Santos, M., Irigoien, X., and Uriarte, A. 2013. Acoustic surveys for juvenile anchovy in the Bay of Biscay: abundance estimate as an indicator of the next year's recruitment and spatial distribution patterns. *ICES Journal of Marine Science*. doi.10.1093/icesjms/fst096.
- Carrera, P. and Porteiro, C. 2003. Stock dynamic of the Iberian sardine (*Sardinapilchardus*, W.) and its implication on the fishery off Galicia. *Scientia Marina.*, 67 (Suppl. 1): 245-258.
- Carrera, P., Churnside, J. H., Boyra, G., Marques, V., Scalabrin, C., and Uriarte, A. 2006. Comparison of airborne lidar with echosounders: a case study in the coastal Atlantic waters of southern Europe. *ICES Journal of Marine Science*, 63: 1736-1750.
- Cochran, W.G. (1977). *Sampling Techniques* (third edition). John Wiley and Sons, New York, 428 pp.
- Coetzee, J. 2000. Use of a shoal analysis and patch estimation system (SHAPES) to characterise sardine schools. *Aquat. Living Resour.*, 13, 1-10.
- Coetzee, J. C., Misund, O. A., and Oechslin, G. 2001. Variable spatial structure of schooling pelagic fish off Namibia: implications for acoustic surveys. *African Journal of Marine Science*, 23 (1) : 99–109.
- De Robertis, A., McKelvey, D. R., and Ressler, P. H. 2010. Development and application of an empirical multifrequency method for backscatter classification. *Canadian Journal of Fisheries and Aquatic Sciences*, 67: 1459–1474.
- Diner, N., Weill, A., Coail, J. Y., and Coudeville, J. M. 1989. INES-MOVIES: a new acoustic data acquisition and processing system. *ICES CM 1989/B:45*. 11pp.
- Dragesund, O. and Olsen, S. 1965. On the possibility of estimating year-class strength by measuring echo-abundance of 0-group fish. *Fiskeridir. Skr. Havundersøk.*, 13:47-75.
- Dressel, S. H., and Norcross, B. L. 2005. Using poststratification to improve abundance estimates from multispecies surveys: a study of juvenile flatfishes. *Fish. Bull.* 103:469-488.
- Edwards, J. I., F. Armstrong. 1984. Herring, mackerel and sprat target strength experiments with behavioural observations. *ICES CM 1984/B:34*. 21 pp.
- Fässler, S. M. M. 2008. target strength variability in atlantic herring (*clupeaharengus*) and its effect on acoustic Abundance estimates. Ph. Thesis. University of St. Andrews. <http://hdl.handle.net/10023/1703>. 277 pp.
- Fässler, S. M. M., O'Donnell, C., and Jech, J.M. 2013. Boarfish (*Caprosaper*) target strength modelled from magnetic resonance imaging (MRI) scans of its swimbladder. *ICES Journal of Marine Science*, 70: 1451–1459.
- Foote, K. G., Aglen, A., and Nakken, O. 1986. Measurement of fish target strength with a split-beam echo sounder. *J. Acoust. Soc. Am.* 80 (2), 612-621

- Foote K.G., 1987. Fish target strengths for use in echo integrator surveys. *J. Acoust. Soc. Am.*, 82: pp. 981-987.
- Frouzova, J., Kubecka, J., Balk, H., and Frouz, J. 2005. Target strength of some European fish species and its dependence on fish body parameters. *Fisheries Research*, 75: 86-89.
- Georgakarakos, S., Trygonis, V., and Haralabous, J. 2011. Accuracy of Acoustic Methods in Fish Stock Assessment Surveys. *Sonar Systems*, Prof. Nikolai Kolev (Ed.), ISBN: 978-953-307-345-3, InTech. Pp 275-298.
- Gorska, N., Korneliussen, R. J., and Ona, E. 2007. Acoustic backscatter by schools of adult Atlantic mackerel. – *ICES Journal of Marine Science*, 64: 1145–1151
- Gutiérrez, M., y MacLennan, D. N., 1998. Resultados preliminares de las mediciones de Fuerza de Blanco in situ de las principales especies pelágicas. *Crucero BIC Humboldt 9803-05 de Tumbes a Tacna*. Inf. Inst. Mar Perú, no. 135:16-19.
- Halldórsson, O. and Reynisson, P. 1983. Target strength measurements of herring and capelin "in situ" at Iceland. *FAO Fish. Rep.* 300: 78-84.
- Hannachi, M.S., Ben Abdallah, L., and Marrakchi, O. 2005. Acoustic identification of small pelagic fish species: target strength analysis and school descriptor classification, *MedSudMed Technical Documents* 5: 90–99.
- Henderson, M. J. 2005. The influence of orientation on the target strength of Pacific hake (*Merluccius productus*). Master Thesis. University of Washington.
- Henderson, M. J., Horne, J. K., and Towler, R. H. 2007. The influence of beam position and swimming direction on fish target strength. *ICES Journal of Marine Science*, 65: 226–237.
- Henderson M. J., and Horne J. K. 2007. Comparison of in situ, ex situ, and backscatter model estimates of Pacific hake (*Merluccius productus*) target strength. *Canadian Journal of Fisheries and Aquatic Sciences*, 64: 1781–1794.
- Huse, I. and Ona, E. 1996. Tilt angle distribution and swimming speed of overwintering Norwegian spring-spawning herring. – *ICES Journal of Marine Science*, 53:863–873.
- Korneliussen, R. J., and Ona, E. 2003. Synthetic echograms generated from the relative frequency response. *ICES Journal of Marine Science*, 60: 636–640.
- Korneliussen, R. J., Diner, N., Ona, E., Berger, L., and Fernandes, P. 2008. Proposals for the collection of multifrequency acoustic data. *ICES Journal of Marine Science*. 65: 982-994.
- Korneliussen, R. J., Heggelund, Y., Eliassen, I. K., and Johansen, G. O. 2009. Acoustic species identification of schooling fish. *ICES Journal of Marine Science*, 66: 1111–1118.
- Lillo S., Cordova J. & Paillaman A., 1996. Target-strength measurements of hake and jack mackerel. *ICES Journal of Marine Science*, 53: pp. 267-272.
- MacCall, A. D. (1990). The dynamic geography of marine fish populations. In *Recruitment fishery oceanography*. (Washington Sea Grant, University of Washington Press, Seattle, 153 pp).
- Machias, A. and Tsimenides, N. Anatomical and physiological factors affecting the swim-bladder cross-section of the sardine *Sardinapilchardus*. *Canadian Journal of Fisheries and Aquatic Science*, 53:280-287.
- Martins, M. M., Skagen, D., Marques, V., Zwolinski, J., and Silva, A. 2013. Changes in the abundance and spatial distribution of the Atlantic chub mackerel (*Scombercolias*) in the pelagic ecosystem and fisheries off Portugal. *Scientia Marina* 77(4) December 2013, 551-563.

- McClatchie, S., Thorne, R. E., Grimes, P. & Hanchet, S. 2000. Ground truth and target identification for fisheries acoustics. *Fisheries Research*, 47, 173-191.
- Muiño, R., Carrera, P., Petitgas, P., Beare, D. J., Georgakarakos, S., Haralambous, J., Iglesias, M., Liorzou, B., Massé, J., and Reid, D. G. 2003. Consistency in the correlation of school parameters across years and stocks. *ICES Journal of Marine Science*, 60: 164–175.
- Muiño, R., Carrera, P., and Iglesias, M. 2003. The characterization of sardine (*SardinapilchardusWalbaum*) schools off the Spanish-Atlantic coast. – *ICES Journal of Marine Science*, 60: 1361–1372.
- Nakken, O. and Dommasnes A. 1975. The application for an echo integration system in investigations on the stock strength of the Barents Sea capelin (*Mallotusvillosus*, Müller) 1971-74. *ICES CM* 1975/B:25.
- Nakken O. & Dommasnes A., 1977. Acoustic estimates of the Barents Sea capelin stock 1971–1976. *ICES CM*, 1977/H:35.
- Ona, E. 1990. Physiological factors causing natural variations in acoustic target strength of fish. *J. mar. biol. Ass. U.K.*, 70: 107-127.
- Patti, B., Mazzola, S.; Calise, L., Bonanno, A., Buscaino, G., and Cosimi, G. 2000. Echo-survey estimates and distribution of small pelagic fish concentrations in the Strait of Sicily during June 1998. *GFCM/ SAC Working Group on Small Pelagics*, Fuengirola, Spain, 1–3 March 2000. 8 pp.
- Pedersen, G., Godø, O. R., Ona, E., and Macaulay, G. J. 2011. A revised target strength–length estimate for blue whiting (*Micromesistiuspoutassou*): implications for biomass estimates. *ICES Journal of Marine Science*, 68: 2222–2228.
- Petitgas, P., Massé, J., Beillois, P., Lebardier, E., and Le Cann, A. 2003. Sampling variance of species identification in fisheries-acoustic surveys based on automated procedures associating acoustic images and trawl hauls. *ICES Journal of Marine Science*, 60: 437-445.
- Pitcher T. J. 1983. Heuristic definitions of shoaling behaviour. *Anim. Behav.*, 31, 611-13.
- Pitcher, T. J. 2001. Fish schooling: implications for pattern in the oceans and impacts on human fisheries. In: *Encyclopedia of Ocean Sciences* (Ed. By J. H. Steele, K. K. Turekian & S. A. Thorpe), pp. 975–987. London: Academic Press.
- Porteiro, C., Carrera, P., and Miquel, J. 1996. Analysis of Spanish acoustic surveys for sardine, 1991-1993: abundance estimates and inter-annual variability. *ICES Journal of Marine Science*, 53: 429-433.
- Reid, D. (ed.) 2000. Report on echo-trace classification. *ICES Cooperative Research Report no 238*, 107 pp.
- Reid, D., Scalabrin, C., Petitgas, P., Massé, J., Aukland, R., Carrera, P., and Georgakarakos, S. 2000. Standard protocols for the analysis of school based data from echosounder surveys. *Fisheries Research*, 47: 125-136.
- Scalabrin, C. 1997. Identification acoustique des espèces pélagiques à partir d'attributs discriminants des bancs de poissons monospécifiques. Thèse du Doctorat en Océanographie Biologique: Université de Bretagne Occidentale de Brest.
- Simmonds, E. J., Williamson, N. J., Gerlotto, F., and Aglen, A. 1992 Acoustic survey design and analysis procedure: a comprehensive review of current practice. *ICES Cooperative Research Report no 187*. 127 pp.

- Simmonds, E. J. & MacLennan, D. N. 2005. Data analysis. In Fisheries Acoustics: Theory and Practice. Oxford: Blackwell.
- Simmonds E. J. and MacLennan, D. 2005. Survey design in Fisheries Acoustics. Theory and practice. 2nd edition. Blackwell Science.
- Simmonds, E.J., Gutiérrez, M., Chipollini, A., Gerlotto, F., Woillez, M., and Bertrand, A. 2009. Optimizing the design of acoustic surveys of Peruvian anchoveta. ICES Journal of Marine Science, 66: 1341–1348.
- Stanton, T. K., Chu, D., Jech, J. M., and Irish, J. D. 2010. New broadband methods for resonance classification and high-resolution imagery of fish with swimbladders using a modified commercial broadband echosounder. – ICES Journal of Marine Science, 67: 365–378.
- Swain, D. P., Sinclair, A.F. 1994. Fish Distribution and Catchability: what is the appropriate measure of distribution?. Can. J. Fish. Aquat. Sci. 51:1046-1054.
- Woillez, M., Poulard, J-C., Rivoirard, J., Petitgas, P., and Bez, N. 2007. Indices for capturing spatial patterns and their evolution in time, with application to European hake (*Merlucciusmerluccius*) in the Bay of Biscay. ICES Journal of Marine Science, 64: 537–550.
- Woillez, M., Rivoirard, J., and Fernandes, P. G. 2009. Evaluating the uncertainty of abundance estimates from acoustic surveys using geostatistical simulations. ICES Journal of Marine Science, 66: 1377–1383.
- Zar, J. H. (1984). Biostatistical Analysis. 2nd edition. Prentice-Hall, Inc.
- Zwolinski, J. 2007. Estimaco acstica da distribuico e abundncia de sardinha, *Sardina pilchardus*. Tesis doctoral. Universidade de Aveiro. Departamento de Biologia. 147pp



IPMA

Instituto Português
do Mar e da Atmosfera